

Somatization, but not depression, is characterized by disorders in the tryptophan catabolite (TRYCAT) pathway, indicating increased indoleamine 2,3-dioxygenase and lowered kynurenine aminotransferase activity

Michael MAES¹, Piotr GALECKI², Robert VERKERK³, Winfried RIEF^{4,5}

¹ Maes Clinics @ Tria, Bangkok, Thailand

² Department of Adult Psychiatry, Medical University of Lodz, Lodz, Poland

³ Department Medical Biochemistry, University of Antwerp, Wilrijk, Belgium

⁴ Klinik Roseneck, Center for Behavioral Medicine, Prien am Chiemsee, Germany

⁵ Department of Clinical Psychology and Psychotherapy, University of Marburg, Marburg, Germany

Correspondence to: Michael Maes, MD., PhD.
Maes Clinics @ TRIA, Piyavate Hospital
998 Rimklongsamsen Road, Bangkok 10310, Thailand.
TEL: +66 26602728; E-MAIL: dr.michaelmaes@hotmail.com

Submitted: 2011-04-01 *Accepted:* 2011-04-22 *Published online:* 2011-06-29

Key words: **tryptophan; serotonin; inflammation; cytokines; oxidative stress; chronic fatigue**

Neuroendocrinol Lett 2011; **32**(3):264–273 PMID: 21712776 NEL320311A13 © 2011 Neuroendocrinology Letters • www.nel.edu

Abstract

BACKGROUND: Reduced plasma tryptophan occurs in depression and somatization. Induction of indoleamine 2,3-dioxygenase (IDO) with consequent synthesis of tryptophan catabolites (TRYCATs) and lowered tryptophan are associated with the onset of depression in the puerperium and during interferon-alpha treatment. Depression is accompanied by lowered kynurenic acid, a neuroprotectant, or increased kynurenine, a neurotoxic TRYCAT.

AIMS AND METHODS: To examine plasma tryptophan; kynurenine; kynurenic acid; the kynurenine / tryptophan (KY/TRP) ratio, indicating IDO activity; and the kynurenine / kynurenic acid (KY/KA) ratio, indicating kynurenine aminotransferase (KAT) activity, in somatization; depression; somatization + depression; and controls. Illness severity is measured by the Somatic Symptom Index (SSI), the Screening for Somatoform Symptoms (SOMS), and the Beck Depression Inventory (BDI).

RESULTS: Tryptophan is significantly lower in patients than in controls and lower in somatization than in depression. KY/TRP is significantly increased in somatization. Kynurenic acid is significantly lower in patients than in controls, and lower in somatization than in depression. KY/KA is significantly higher in somatization and somatization + depression than in depression and controls. There are significant correlations between the severity of somatization, but not depression, and KY/TRP and KY/KA (positive) and tryptophan (negative). Kynurenine and kynurenic acid are significantly correlated in controls, somatization + depression, and depression, but not in somatization.

CONCLUSIONS: Somatization is characterized by increased IDO activity and disorders in KAT activity and an increased neurotoxic potential. The TRYCAT pathway may play a role in the pathophysiology of somatizing and “psychosomatic”

symptoms through effects on pain, gut motility, the autonomic nervous system, peripheral NMDA receptors, etc. Even more, biological disorders, such as aberrations in the TRYCAT pathway, which are considered to be a hallmark for depression, are in fact attributable to somatization rather than to depression per se. Future research in depression on the TRYCAT pathway should always control for the possible effects of somatization.

INTRODUCTION

There is evidence that depletion of plasma tryptophan, and induction of indoleamine 2,3-dioxygenase (IDO) with consequent synthesis of tryptophan catabolites (TRYCATs) play a role in the onset of depression (Maes et al. 2011). The conversion of tryptophan to kynurenine is the first and rate-limiting step in this pathway. Kynurenine is further metabolized into kynurenic acid by kynurenine 2,3-aminotransferase (KAT) or to nicotinamide. IDO is widely expressed in human tissues, e.g. the brain, kidney, lung, spleen, and duodenum, macrophages and dendritic cells (Takikawa et al. 1984; Moroni et al. 1991). IDO is induced primarily by interferon- γ (IFN γ) and cytokines, like interleukin-1 β (IL-1 β) and tumor necrosis factor- α (TNF α) (Maes et al. 2011; Oxenkrug 2007). There is evidence that depression is accompanied by cell-mediated immune activation, as indicated by increased IFN γ and neopterin; and an inflammatory response, with increased production of IL-1 β and TNF α (Maes 1995; 2010; Maes et al. 2011).

Activation of IDO diverts tryptophan from the 5-HT synthetic route thereby depleting tryptophan and consequently brain 5-HT and leads to the formation of some TRYCATs that are important to depression, e.g. kynurenine, and kynurenic acid (Maes et al. 1994; 2011). Indeed, kynurenine may be detrimental as it has depressogenic, anxiogenic, excitotoxic and neurotoxic effects. Kynurenic acid, on the other hand, has neuroprotective effects. Therefore, the kynurenine / kynurenic acid (KY/KA) ratio indicates not only KAT activity, but also the neurotoxic potential generated during IDO activation.

Depletion of plasma tryptophan is frequently observed in depression (DeMyer et al. 1981; Joseph et al. 1984; Moller 1985; Maes et al. 1987a; 1991a; Cowen et al. 1989). Low plasma tryptophan is significantly associated to various markers of cell-mediated immune activation in depression (Maes et al. 1993; 1994; Song et al. 1998). This indicates that cell-mediated immune activation and the inflammatory response in depression have induced IDO thus causing lowered plasma tryptophan. In the early puerperium, IDO activation, as estimated by the kynurenine/tryptophan (KY/TRP) ratio, and TRYCAT production are significantly associated with the development of affective symptoms (Maes et al. 2001b; 2002). The onset of affective symptoms during IFN α -based immunotherapy is significantly related to the KY/TRP ratio, plasma kynurenine and the KY/KA ratio (Maes et al. 2001a; Bonaccorso et al. 2002; Wichers et al. 2005b). Raison *et al.* (2010) reported that

the IFN α -induced synthesis of kynurenine and quinolinic acid, another neurotoxic TRYCAT, are associated with the onset of depression. In depressed patients there are only few data: some authors found increased levels of neurotoxic TRYCATs (Gabbay et al. 2010), whereas others found lowered levels of neuroprotective TRYCATs (Myint et al. 2007).

Depression shows a high degree of comorbidity with somatization, a multi-symptomatic syndrome characterized by unexplained physical symptoms, that has a prevalence rate of around 4–7% in the general population (Escobar et al. 1987; Rief et al. 2001a). Other terms that are frequently used to describe this prevalent disorder are 'functional somatic symptoms', 'psychosomatic syndrome' or 'somatoform disorders' (Rief et al. 2004). There is a considerable overlap between somatization and chronic fatigue syndrome, an illness that shows a strong comorbidity with depression (Rief et al. 2010). In primary care, depression is the most common comorbid diagnosis (54.6%) of somatization disorder (Brown et al. 1990). In the NIMH Epidemiologic Catchment Area study (Simon & vonKorff 1991), the increasing number of symptoms in subjects with somatization was strongly associated with depression. Recently, we have reported that patients with somatization have a lowered availability of plasma tryptophan and signs of monocytic activation, such as increased soluble IL-1 receptor antagonist (sIL-1RA) levels, and suppressed cell-mediated immunity, as indicated by lowered levels of the sCD8 molecule (Rief et al. 2004; 2001b). These results show that monocytic activation may underlie the pathophysiology of somatization. Likewise in related disorders, such as chronic fatigue syndrome and fibromyalgia, signs of inflammation and oxidative stress have been observed (Maes & Twisk, 2010; Maes et al. 2011; Brkic et al. 2010).

There are, however, no data whether the lowered plasma tryptophan in somatization may be explained by IDO activation or whether other disturbances in the TRYCAT pathway are involved in the pathophysiology of somatization.

This study was carried out to examine: a) whether increased IDO activation may explain lowered tryptophan levels in depression and somatization; b) whether depression is characterized by an increased neurotoxic (kynurenine) potential and/or decreased neuroprotective (kynurenic acid) potential; and c) other disorders in the TRYCAT pathway in somatization disorder.

SUBJECTS AND METHODS

Subjects

One-hundred and forty-six subjects participated in this study: one hundred and eleven patients and 35 normal controls. All patients were admitted to the Roseneck Center for Behavioral Medicine, a German inpatient treatment Center for psychiatric and psychosomatic disorders. Before inclusion in the study, patients were examined by means of self-rating scales, i.e. Beck

Depression Inventory (BDI) (Beck et al. 1961) and the Screening for Somatoform Symptoms (SOMS) (Rief et al. 1997). The SOMS is a self-rating scale that considers the presence of 53 physical symptoms during the last two years. The symptoms of the SOMS include all somatoform symptoms mentioned in the classification criteria of DSM-IV somatization disorder, symptoms of the ICD-10 somatization disorder and the ICD-10 somatoform autonomic dysfunction list. Only patients with eight or more unexplained physical symptoms on the SOMS or a BDI-score of 17 or above were allowed to participate in the study. Diagnoses were made using a standardized psychiatric interview carried out by senior psychiatrists or clinical psychologists. We used the German translation of the Structured Clinical Interview for DSM-IV SCID (Wittchen et al. 1997) to assess present and lifetime diagnoses of major depression, somatization disorder, and other psychiatric disorders. In order to facilitate the diagnostic procedures we used the International Diagnostic Checklist – IDCL (Hiller et al. 1997).

Depression was diagnosed according to the DSM-IV criteria, using the SCID interview, on preselected patients who had a BDI-score of 17 or above. The diagnosis 'somatization' was based on the DSM-IV criteria for somatization disorder but defined as a multisomatiform disorder with a cut-off score of 8 symptoms or more (SSI-8) on the Somatization Index (SSI). The SSI is thus based on a structured interview based on the DSM-IV criteria for somatization disorder that includes a list of 33 medically unexplained physical symptoms. Our diagnosis of somatizing thus constitutes a subthreshold syndrome of the DSM-IV somatization disorder. In our experience, somatization is a more relevant diagnostic entity than somatization disorder. The criteria for somatization disorder are over-exclusive and do not represent the healthcare relevance of patients with multiple somatoform symptoms (Escobar et al. 1989). Previously, we have externally validated the diagnosis "somatization" because we have shown that the SSI-8 criterion constitutes a useful index to differentiate patients with high versus low disabilities (Rief et al. 1997). Comparable results have been published using the diagnosis "multisomatiform disorder", defined as 3 or more medically unexplained symptoms plus a long history of somatization (> or = 2 years) (Kroenke et al. 1997). Finally, we included 36 patients with somatization (SOM), 36 patients with depression (MDD), 39 patients with 'comorbid' SOM and MDD (SOM+MDD). Nine of the 36 SOM patients fulfilled the criteria for somatization disorder, while 15 of the 38 patients with SOM+MDD were classified as DSM-IV somatization disorder. Patients and controls gave written informed consent after the study protocol was fully explained; the study has been approved by the local ethical committee.

We excluded subjects with any medical illnesses, e.g. inflammatory bowel disorders, diabetes, hypertension, cardiovascular disorders and any condition that could

have explained the somatoform symptoms; b) subjects with acute inflammatory or allergic reactions the last 2 months prior to the study; c) patients with life-time diagnoses of other axis-I DSM IV disorders, e.g. psychotic, substance use and organic mental disorders; and d) subjects who had been treated with immunomodulatory drugs, such as glucocorticoids, etc. Only subjects who were free of psychotropic drugs, including antidepressants, were selected to participate in this study. Thirteen women took contraceptive drugs, 8 subjects took thyroxine, and some were taking antioxidants. We have controlled our results by using the following variables as explanatory variables in regression analyses: drug state (drug free versus not drug free), smoking behavior, alcohol use and use of caffeine, and the body mass index (BMI), as calculated as weight (kg) / body height (in meter)².

Severity of depression was measured by means of Beck Depression Inventory (BDI) (Beck et al. 1961). The 90-item version of the Symptom Check List-90-Revised (SCL-90) (Derogatis 1994; Rief & Fichter 1992) was employed to measure the general severity of psychopathology according to different symptom dimensions. The SSI was used as a general index for the severity of somatization. The DSM-IV SSI-index is the total score of positively answered symptoms out of 33 bodily complaints. The retest reliability of the SSI is $r_{tt} = 0.85$ (72 h). Self-rated severity of somatic symptoms was assessed by means of the SOMS (Rief et al. 1997). The SSI and the SOMS showed a strong correlation, i.e. $r = 0.75$.

Methods

After an overnight fast, blood (2×7 cc) was sampled at 7:00 a. m. for the assay of the biomarkers. Subjects were instructed to stay in bed until the blood sample was taken. At 8:00 a. m. blood samples were centrifuged at 1500 g for 10 min at 4°C and stored at -70°C until thawed for assay. The initial interview, the self-rating scales and the blood samplings were carried within a time window of 3–4 days. Plasma specimens for the assay of tryptophan, kynurenine, and kynurenic acid were assayed in a single run with a single lot number of reagents and consumables employed by a single operator (who was blind to the diagnosis) in order to minimize variation in the results. Tryptophan was assayed by means of high performance liquid chromatography (HPLC) as previously explained (1996). Serum kynurenine and kynurenic acid were determined by means of HPLC as described by Herve *et al.* (1996). Both TRYCATs were analyzed in deproteinised plasma samples employing reversed phase HPLC with Chromolith performance PR-18e, 4.7 100 mm column (Merck, Darmstadt, Germany). Kynurenine was detected spectrophotometrically at 365 nm and kynurenic acid fluorimetrically at 334 nm and an emission wavelength of 388 nm. The mobile phase was prepared with 250 mM zinc-acetate in distilled water (27.4 g in 500 ml) and pH was brought to 5.8 with acetic acid and made up to a

volume of 455 ml with water. A total of 45 ml acetonitrile was added into this mixture. We prepared the kynurenine standards by dissolving kynurenine in water and the kynurenic acid standards by dissolving kynurenic acid in ethanol and hydrochloric acid. For the working standards, stock standards of 5 μ M for kynurenine (500 μ l) and 100 nM for kynurenic acid (10.0 μ l) were diluted in distilled water. Perchloric acid was used for deproteinization. Tryptophan and kynurenine are expressed as μ mol/L and kynurenic acid as nmol/L. The intra- and interassay CV values for all analytes obtained in our laboratory were less than 5% and 7%, respectively. The kynurenine/tryptophan quotient (KY/TRP) was computed, since this ratio allows to estimate IDO activity. The kynurenic acid / kynurenine (KY/KA) quotient was computed as an index for neurotoxic/neuroprotective potential and KAT activity.

Statistics

Group mean differences were examined by means of analysis of variance (ANOVA) and analysis of covariance (ANCOVA). Multiple comparisons between group means were assessed with the Dunn test. Relationships between variables were checked by means of Pearson's product moment correlation coefficients, partial correlation coefficients, multiple regression analyses and canonical correlation analyses. The significance of the difference between two correlation coefficients between the same sets of variables in two independent study groups was calculated employing a Fisher r-to-z transformation. The independence of classification systems was checked by means of analysis of contingency (χ^2 -test). Transformations (Box-Cox) were used to normalize the distribution of the data if necessary. The level of significance was set at $p=0.05$ (two-tailed).

RESULTS

Demographic data

Table 1 shows age, gender ratio, the SSI, SOMS, BDI, SCL-90, and the biomarkers in patients with SOM, SOM+MDD, MDD, and normal controls. There were no significant differences in age ($F=1.4$, $df=3/142$, $p=0.2$) and gender ratio ($\chi^2=4.1$, $df=3$, $p=0.2$) between the 4 study groups. The SSI was significantly higher in SOM+MDD than in SOM ($t=3.76$, $p=0.0005$), MDD ($t=14.34$, $p<10^{-5}$) and controls ($t=19.32$, $p<10^{-5}$); higher in SOM than in MDD ($t=10.37$, $p<10^{-5}$) and controls ($t=15.28$, $p<10^{-5}$); and higher in MDD than in controls ($t=4.99$, $p=0.00003$). The same pattern was observed for the SOMS, BDI and SCL-90, except that the differences in the SOMS between SOM+MDD and SOM were less significant ($t=2.40$, $p=0.016$); the BDI ($t=1.52$, $p=0.12$) and SCL-90 ($t=1.73$, $p=0.08$) were not significantly different between MDD and MDD+SOM.

In the total study group, there were significant correlations between plasma tryptophan and kynurenine ($r=0.45$, $p<10^{-5}$) and kynurenic acid ($r=0.30$, $p=0.0005$).

There was a significant and positive correlation between kynurenine and kynurenic acid ($r=0.50$, $p<10^{-5}$). There was no significant correlation between the KY/TRP and KY/KA ratios ($r=0.09$, $p=0.3$). The effects of age and sex on the biomarkers were examined by means of multiple regression analysis. Age ($F=17.6$, $p=0.0002$) and sex ($F=14.8$, $p=0.0004$) explained 17.1% of the variance in plasma tryptophan ($F=15.7$, $df=2/141$, $p=0.00002$). Sex ($F=9.0$, $p=0.003$), but not age ($F=0.7$, $p=0.6$) explained 6.5% of the variance in plasma kynurenine ($F=4.9$, $df=2/141$, $p=0.008$). Sex ($F=8.4$, $p=0.004$), but not age ($F=3.5$, $p=0.006$) explained 7.9% of the variance in plasma kynurenic acid ($F=6.2$, $df=2/141$, $p=0.003$). Age ($F=22.1$, $p=0.00005$) and sex ($F=0.3$, $p=0.6$) explained 12.4% of the variance in the KY/TRP ratio ($F=11.1$, $df=2/141$, $p=0.0001$). Age ($F=4.4$, $p=0.036$) and sex ($F=3.9$, $p=0.046$) explained 5.7% of the variance in the KY/KA ratio ($F=4.3$, $df=2/141$, $p=0.01$). Thus, age had a significant effect on tryptophan (decreasing), the KY/TRP ratio (increasing) and the KY/KA ratio (decreasing). Gender has a significant effect on tryptophan, kynurenine and kynurenic acid (higher in men) and the KY/KA ratio (higher in women). Therefore, we have adjusted the following analyses for possible effects of age and sex by entering these as additional explanatory variables in ANCOVAs or partial correlation coefficients.

Intergroup differences

ANCOVA with age as covariate and sex as second factor showed significant differences in plasma tryptophan between the 4 groups ($F=17.7$, $df=3/137$, $p<10^{-5}$). Dunn tests showed that plasma tryptophan was significantly lower in SOM as compared to SOM+MDD ($t=2.25$, $p=0.02$), MDD ($t=3.68$, $p=0.0006$) and the controls ($t=7.61$, $p<10^{-5}$). There were no significant differences in plasma tryptophan between SOM+MDD and MDD ($t=1.50$, $p=0.1$). SOM+MDD ($t=5.52$, $p=0.00001$) and MDD ($t=3.95$, $p=0.0003$) patients had lower plasma tryptophan than the controls.

ANCOVA (Table 1) showed significant differences in plasma kynurenine between the groups ($F=3.6$, $df=3/136$, $p=0.01$). Dunn tests showed that patients with SOM ($t=3.22$, $p=0.002$) and SOM+MDD ($t=2.98$, $p=0.003$) had significantly lower kynurenine than controls. An ANCOVA with plasma tryptophan as additional explanatory variable showed that the intergroup differences in plasma kynurenine disappeared ($F=0.2$, $df=3/135$, $p=0.9$) after adjusting for tryptophan ($F=17.5$, $df=1/135$, $p=0.0002$).

ANCOVA (Table 1) shows significant differences in plasma kynurenic acid between the groups ($F=9.3$, $df=3/137$, $p=0.00006$). Kynurenic acid was significantly lower in SOM and SOM+MDD than in MDD ($t=3.22$, $p=0.002$, and $t=2.77$, $p=0.006$, respectively) and controls ($t=5.21$, $p=0.00002$ and $t=4.80$, $p=0.00004$, respectively). There were no significant differences in kynurenic acid between SOM and SOM+MDD ($t=0.52$,

$p=0.61$). Kynurenic acid is significantly lower in MDD as compared to controls ($t=2.01$, $p=0.043$).

ANCOVA (Table 1) shows significant differences in the KY/TRP ratio between the diagnostic groups ($F=2.75$, $df=3/139$, $p=0.04$). Dunn tests shows a significantly higher KY/TRP ratio in SOM than in SOM+MDD ($t=1.96$, $p=0.048$) and controls ($t=3.59$, $p=0.0007$), but no significant differences between SOM and MDD ($t=1.83$, $p=0.07$) and SOM+MDD ($t=1.69$, $p=0.09$) or MDD ($t=1.75$, $p=0.08$) versus normal controls.

ANCOVA (Table 1) shows significant differences in the KY/KA ratio between the diagnostic groups ($F=8.6$, $df=3/140$, $p=0.0001$). Dunn tests shows a significantly higher KY/KA ratio in SOM than in MDD ($t=3.02$, $p=0.003$) and controls ($t=4.52$, $p=0.00008$); a significantly higher KY/KA ratio in SOM+MDD than in MDD ($t=2.32$, $p=0.02$) and controls ($t=3.86$, $p=0.0004$); and no significant differences between MDD and controls ($t=1.54$, $p=0.1$). All above results of ANCOVAs were not affected by entering the drug state, smoking behavior, alcohol use, use of caffeine, or BMI.

Correlations between tryptophan and the TRYCATs

As described above, there are significant correlations between tryptophan and kynurenine, while ANCOVA showed that the KY/TRP ratio is increased in SOM patients only; there are also correlations between kynurenine and kynurenic acid, while ANCOVA shows that the KY/KA ratio is increased in patients with SOM and SOM+MDD only. This suggests that the rates of both reactions, which are dependent on the levels of IDO and tryptophan, or on KAT and kynurenine may be different between the groups. Therefore, we have computed the correlations between tryptophan and kynurenine, and between kynurenine and kynurenic

acid in the 4 study groups separately. Tryptophan and kynurenine were significantly correlated in the controls ($r=0.39$, $p=0.018$), SOM+MDD ($r=0.37$, $p=0.02$) and MDD ($r=0.45$, $p=0.007$), but not in SOM ($r=0.14$, $p=0.6$). By means of the Fisher r -to- z transformation no significant differences could be detected between those correlations coefficients. Kynurenine and kynurenic acid were significantly correlated in the controls ($r=0.58$, $p=0.0004$), SOM+MDD ($r=0.54$, $p=0.0007$) and MDD ($r=0.53$, $p=0.001$), but not in SOM ($r=0.06$, $p=0.7$). Using the Fisher r -to- z transformation, significant differences were found between the correlation coefficient obtained in SOM and those in normal controls ($z=-2.43$, $p=0.007$); SOM+MDD ($z=-2.26$, $p=0.01$); and MDD ($z=-2.15$, $p=0.016$).

Correlations with severity of illness

In the total study group there were significant correlations between the SSI and the SOMS ($r=0.80$, $p<10^{-5}$), BDI ($r=0.52$, $p<10^{-5}$), and SCL-90 ($r=0.49$, $p<10^{-5}$); the SOMS and the BDI ($r=0.62$, $p<10^{-5}$) and SCL-90 ($r=0.70$, $p<10^{-5}$); and between the BDI and the SCL-90 ($r=0.74$, $p<10^{-5}$). Semipartial correlation analyses (after adjusting for sex and age) showed significant relationships between tryptophan and the SSI ($r=-0.36$, $p=0.00005$), the SOMS ($r=-0.36$, $p=0.00007$), but not the BDI ($r=-0.16$, $p=0.053$) and the SCL-90 ($r=-0.17$, 0.051); between kynurenine and the SSI ($r=-0.16$, $p=0.047$), and the SOMS ($r=-0.23$, $p=0.006$), but not the BDI ($r=-0.10$, $p=0.2$) and the SCL-90 ($r=-0.07$, 0.6); between kynurenic acid and the SSI ($r=-0.26$, $p=0.002$), the SOMS ($r=-0.36$, $p=0.00008$), the BDI ($r=-0.18$, $p=0.03$) and the SCL-90 ($r=-0.19$, 0.03); between the KY/TRP ratio and the SSI ($r=0.18$, $p=0.03$), but not the SOMS ($r=0.10$, $p=0.2$), the BDI ($r=0.04$, $p=0.6$) and the SCL-90 ($r=0.08$, 0.6); and between the KY/KA

Tab. 1. Demographic data, rating scales and measurements of plasma tryptophan, kynurenine, kynurenic acid and the kynurenine/tryptophan (KY/TRP) and kynurenine/kynurenic acid (KY/KA) ratio's in patients with somatization (SOM), SOM and depression (MDD), MDD and in normal controls.

Variables	SOM	SOM+MDD	MDD	controls
Age (years)	43.1 (12.0)	40.2 (10.4)	41.9 (9.9)	38.0 (11.9)
Gender ratio (M/F)	13 / 23	10 / 29	13 / 23	17 / 18
SSI	9.5 (2.3)	11.6 (3.3)	3.7 (2.3)	0.9 (1.0)
SOMS	19.0 (5.9)	22.2 (6.5)	12.8 (6.6)	2.4 (2.5)
BDI	13.9 (5.3)	29.2 (7.5)	27.0 (8.2)	2.7 (3.1)
SCL-90	50.1 (16.3)	64.9 (10.8)	59.4 (14.6)	16.4 (12.6)
Tryptophan ($\mu\text{M/L}$)	58.9 (8.3)	65.8 (10.7)	69.2 (14.6)	81.1 (14.0)
Kynurenine ($\mu\text{M/L}$)	1.67 (0.30)	1.69 (0.36)	1.80 (0.48)	1.96 (0.42)
Kynurenic acid (nmol/L)	20.95 (11.53)	22.29 (13.59)	28.50 (15.46)	31.80 (11.03)
KY/TRP $\times 100$	2.87 (0.63)	2.59 (0.54)	2.62 (0.64)	2.44 (0.51)
KY/KA	95 (38)	92 (33)	75 (25)	67 (20)

SSI: Somatization Index; SOMS: Screening for Somatoform Symptoms; BDI: Beck Depression Inventory; SCL-90: Symptom Check List-90-Revised; All results are shown as mean (SD). For the results of the statistical analyses: see text

ratio and the SSI ($r=0.23$, $p=0.005$), the SOMS ($r=0.30$, $p=0.0005$), but not the BDI ($r=0.15$, $p=0.06$) and the SCL-90 ($r=0.17$, $p=0.052$).

Table 2 shows the outcome of a canonical correlation analyses whereby the 4 rating scales were entered as dependent variables and plasma tryptophan, the KY/TRP ratio and the KY/KA ratio as independent variables. Two significant eigenvectors were found: the first ($r=0.43$, $p<0.01$) showed high loadings on the SSI, the SOMS, the SCL-90 (positively loaded), and the KY/KA ratio (positively loaded) and tryptophan (negatively loaded); whereas the second ($r=0.21$, $p<0.05$) loaded highly on the SSI and the KY/TRP ratio.

DISCUSSION

The first major finding of this study is that, although plasma tryptophan was significantly lower in depressed patients, IDO activity was not increased. These results are in contrast to our primary hypothesis and to previous studies in women in the puerperium and IFN α -treated patients. In these studies, a significant association between increased IDO activity and affective symptoms was detected (Maes et al. 2002; 2001b; Bonaccorso et al. 2002; Wichers et al. 2005b; Raison et al. 2010). Also, Gabbay et al. (2010) found a significantly higher KY/TRP ratio and lower tryptophan in adolescent depression with melancholic features as compared to non-melancholic depressed adolescents and controls. In patients with coronary artery disease, an elevated KY/TRP ratio was significantly correlated with increased depression scores (Swardfager et al. 2009). In malignant diseases, IDO activation and lowered plasma tryptophan parallel the course of the disease and are related to the onset of depressive mood (Brandacher et al. 2006). However, also Myint et al. (2007) could not find a significant difference in the KY/TRP ratio between depressed patients and controls.

There are other mechanisms in depression that can explain lowered plasma tryptophan even in the absence of IDO activation. A first possible mechanism is enhanced catabolism of tryptophan along the TRYCAT pathway by activation of liver tryptophan 2,3-dioxygenase (TDO, tryptophan pyrolase EC 1.13.11.11) by glucocorticoids, which are known to be increased in depression (Maes et al. 2011). In this respect, it has been shown that dexamethasone or glucocorticoid administration in rats enhances liver TDO activity and consequently lowers tryptophan in the liver, serum and brain, as well as 5-HT in the brain (Morgan & Badawy 1989; Young 1981). TDO knocked-out mice show elevated plasma tryptophan and tryptophan and 5-HT levels in the hippocampus and midbrain (Kanai et al. 2009). In humans, administration of 1 mg dexamethasone is sufficient to lower plasma tryptophan in controls and depressed patients (Maes et al. 1990a; 1990c). Another inflammatory mechanism that may explain lowered plasma tryptophan in depression is related to serum

Tab. 2. Results of canonical correlation analysis performed on both the clinical severity ratings, as the first set of variables, and plasma tryptophan, the kynurenine/tryptophan (KY/TRP) and kynurenine/kynurenic acid (KY/KA) ratio, as a second set of variables.

variables	first canonical eigenvector	second canonical eigenvector
SSI	0.831	-0.352
SOMS	0.938	0.175
BDI	0.274	0.211
SCL-90	0.507	-0.119
Tryptophan	-0.856	0.300
KY/TRP ratio	0.243	-0.964
KY/KA ratio	0.585	0.079
canonical correlation coefficient	$r=0.43$ ($p<0.01$)	$r=0.21$ ($p<0.05$)

The significant loadings (>0.350) are shown in bold. Tryptophan, the KY/TRP and KY/KA ratios were entered as the residualized values after adjusting for age and gender.

albumin, one of the negative acute phase proteins, that is reduced in depression (Maes et al. 1991b; Song et al. 1994). Total plasma tryptophan consists of a major fraction, i.e. 70–90%, that is loosely bound to albumin and a smaller fraction that circulates as free tryptophan (Curzon & Sarna 1984; Fernstrom 1984; Yuwiler et al. 1977). The significant and positive correlations that are detected in depression between plasma tryptophan and serum albumin indicate that the reduced serum albumin levels may have depleted plasma total tryptophan (Maes et al. 1996; 1997). Finally, the failure to find increased IDO activity in our depressed patients does not mean that the lowered availability of tryptophan in depression is not the consequence of IDO activation. Indeed, previously we have reviewed that activation of IDO may, through reduced plasma tryptophan and increased TRYCAT levels, exert a negative feedback on the primary immune-inflammatory response in depression thereby acting as a counter anti-inflammatory response system (CARS) (Maes et al. 2007; 2011; Burdette et al. 2010). Tryptophan depletion leads to metabolic shutdown and starvation of immune cells, whereas increased TRYCATs lead to T cell unresponsiveness and anti-inflammatory effects, which in turn attenuate IDO activation (Maes et al. 2007; 2011). Thus, subchronic conditions with an activated CARS may be characterized by a normalized KY/TRP ratio and lowered plasma tryptophan levels. Even more, in our study we found that after covarying for tryptophan in an ANCOVA the differences in kynurenine between depressed patients and controls disappeared. This indicates that the lowered plasma levels of kynurenine in depression are a consequence of depletion of its precursor, tryptophan.

The second major finding of this study is that patients with somatization had an increased KY/TRP ratio, indicating increased IDO activity. Therefore, the lowered plasma tryptophan in somatization may be explained, at least in part, by activation of IDO. We

should underscore that the plasma levels of tryptophan are much more reduced in somatization than in depressed patients (this study) and more reduced than for example during treatment with IFN α -based immunotherapy (Bonaccorso et al. 2002). In the present study we found that the lowered tryptophan was significantly related to the SSI and the SOMS, but not the BDI and the SCL-90. Also the KY/TRP ratio was significantly associated with the SSI but not the SOMS, BDI or SCL-90. This indicates that lower tryptophan and activated IDO are biomarkers for somatization rather than for depression. Previously, it was found that in depression, lower plasma and CSF tryptophan are associated with anxiety and somatization (Joseph et al. 1984), agitation (Curzon 1979), neuromuscular symptoms, anxiety, agitation, depressed mood and catatonia (Lehmann 1972), and psychic anxiety, depersonalization, obsessions, paranoid symptoms and diurnal variation (Maes et al. 1990b). IDO activation following IFN α -based immunotherapy is also associated to the onset of somatic symptoms, which develop soon after starting treatment and which predict the outcome of the more cognitive symptoms of depression (Wichers et al. 2005a; 2005b). Terre et al. (2003) reported that somatic complaints may represent one risk factor for the subsequent development of depression. The activation of IDO in somatization may offer an explanation for the gender-specific rate of somatization and somatization disorder with a 1.6–2/1 and 10/1 female/male prevalence rate, respectively (Ladwig et al. 2001; Swartz et al. 1990). Thus, women show significantly greater abnormal responses in the TRYCAT pathway than their male counterparts. For example, depressed women excrete significantly more xanthurenic acid, another TRYCAT, following 5g tryptophan than male patients (Maes et al. 1987b). Plasma tryptophan and the KY/TRP ratio following IFN α -based immunotherapy are significantly more depleted and enhanced, respectively, in women than in men (Maes et al. 2011; Bonaccorso et al. 2002). This indicates that IDO is more vulnerable to immune-inflammatory challenges in women and that these gender-related differences in IDO activation could play a role in the increased prevalence of somatization in females.

The third major finding of this study revolves around the concentrations of the detrimental TRYCAT kynurenine versus that of the neuroprotective TRYCAT kynurenic acid, which were normal versus reduced in depression. First, the negative kynurenine findings are in accordance with those of Myint et al. (2007) who found no significant differences in kynurenine concentrations between depressed patients and controls. These negative findings contrast the results that in the puerperium and during IFN α -based immunotherapy the production rate of kynurenine is associated to the onset of depressive symptoms (Maes et al. 2001a; Bonaccorso et al. 2002; Raison et al. 2010). In depressed adolescents, Gabbay et al. (2010) found significant increases in kynurenine in adolescents with melancholic features, while there

was a significant correlation between kynurenine levels and severity of depression. Thus, differences in population, i.e. acute (puerperium and IFN α -based treatment) versus subchronic (clinical depression) conditions, and melancholic adolescents versus depressive adults, could explain the differences between these studies. Second, as Myint et al. (2007) we found lowered kynurenic acid levels in depression, whereas we were unable, in contrast to Myint et al. to find an increased KY/KA ratio in depression (or as they express it a lowered KA/KY index). In Myint's study, however, the results were not adjusted for the significant effects of age on the KY/KA ratio by means of regression analysis. There is another major difference between our study and all above-mentioned studies which may explain differences, i.e. we used a selected group of depressed patients without concomitant 'somatization', whereas all previous researchers did not take somatization into account. As we explained and will explain in the next paragraph, somatization as defined in our study is of critical importance.

So, the fourth major finding of this study is that patients with somatization and comorbid somatization and depression had significantly lowered kynurenic acid levels than depressed patients and normal controls. Likewise, the KY/KA index was significantly increased in both study groups as compared to controls. Correlation analyses revealed significant associations between the KY/KA ratio and the SSI and the SOMS, but not the BDI or the SCL-90. This shows that disorders in the TRYCAT pathway are in fact specific for somatization and not for depression per se and that disorders in the TRYCAT pathway can be found in depressed samples when somatization is present.

The fifth major finding of our study is that there are significant differences in the correlation coefficients between kynurenine and kynurenic acid obtained in somatization ($r=0.06$, $p=0.7$) versus those in normal controls ($r=0.58$, $p=0.0004$), comorbid somatization and depression ($r=0.54$, $p=0.0007$) and depression ($r=0.53$, $p=0.001$). This suggests that the rate of product formation (reaction rate of kynurenic acid) from its substrate (kynurenine), which depends on KAT levels, differs considerably between somatization and all other study groups. These findings and the lowered product (kynurenic acid) levels in somatization suggest that the KAT enzyme concentrations are reduced or that the properties of KAT (K_m , k_{cat}) are modified in somatization. This may be caused by inhibitory effects by endogenous KAT inhibitors, or through modifications of KAT's properties by post-translational modifications or gene mutations.

The lowered kynurenic acid and thus increased KY/KA ratio, pointing toward a relative kynurenine excess in somatization, and the lower tryptophan availability may play a role in symptom formation in somatization patients by modulating pain, anxiety, the autonomic nervous system, gut motility, peripheral nerve func-

tion, ventilation, cardiac functions, etc. Thus, kynurenine contributes to pain and gut motility (Stone & Darlington 2002). Kynurenic acid, on the other hand, has significant antinociceptive and analgesic effects and inhibits intestinal hypermotility (Mecs et al. 2009; Kekesi & Horwath 2002; Kaszaki et al. 2008). Kynurenine is neurotoxic for the peripheral nerves, whereby an excess of serum kynurenine is associated with the development of peripheral neuropathy (Huengsborg et al. 1998). Moreover, peripheral N-methyl-D-aspartate receptors (NMDARs) are detected in deep tissues and are involved in deep tissue pain (Cairns et al. 2003).

Injection of glutamate into the masseter muscle provokes afferent discharges in rats and muscle pain in humans through activation of peripheral NMDARs (Cairns et al. 2003). The same authors concluded that peripherally acting NMDAR antagonists may constitute effective analgesics for NMDA-related muscle pains. Peripheral NMDARs are now also regarded as novel targets for treatment of neuropathic pain (Wu & Zhuo 2009). NMDARs expressed on spinal afferent neurons are upregulated in the lumbosacral dorsal root ganglia following experimental colitis (Li et al. 2006). Peripheral NMDARs play a role in behavioral pain responses to colonic distention in the gut, suggesting that these receptors are important in visceral pain transmission (McRoberts et al. 2001). The glutamate-induced activation of the NMDARs is regulated by TRYCATs. Kynurenic acid is the only endogenous antagonist of NMDARs and acts as an inhibitor of glutamate release (Nemeth et al. 2005). Thus, a lowered KAT activity or kynurenic acid levels – as detected in somatization – could be involved in the maintenance of persistent pain-related behaviors. Important is that peripheral inflammation – as detected in somatization (Rief et al. 2001) – could have increased the expression of those peripheral NMDA receptors that lead to behavioral sensitization during inflammatory pain (Yang et al. 2009).

Not only KAT activity but also the depletion of plasma tryptophan may play a role in the onset of somatizing symptoms. It is well established that changes in the availability of plasma tryptophan determine the synthesis of 5-HT in the brain (Moir & Eccleston 1968). There is also evidence that tryptophan depletion techniques [for example through administration of tryptophan-free amino acid drinks coupled with ingestion of large concentrations of amino acids that compete for the same amino acid transporter] induce depressive symptoms in some patients who had suffered from depression (Maes & Meltzer 1995). Tryptophan depletion also provokes somatization symptoms in humans, e.g. increased visceral perception, and increased pain and higher urge scores during rectal distention in patients with irritable bowel syndrome (Kilkens et al. 2004); increased basal ventilation (Struzik et al. 2002); increased nausea, headache, light-induced pain and photophobia in migraine patients (Drummond 2006); lowered heart rate in subjects with attention deficit disorder (Zepf et al. 2009); increased systolic and diastolic blood pressure and psychological responses to

stress (Davies et al. 2006), and decreased heart rate variability (Booij et al. 2006); increased autonomic stress responses in patients with social anxiety (van Veen et al. 2009); increased distress following uncontrollable stressors in humans (Richell et al. 2005) and stress-sensitivity in rats (Tanke et al. 2008). Behavioral correlates of 5-HT depletion include enhanced pain responses (Harvey et al. 1975).

This study and our previous studies (Rief et al. 2001) show that the unexplained physical symptoms in somatization may in fact have a biological explanation. Even more, biological disorders, such as aberrations in the TRYCAT pathway, which are considered to be a hallmark for depression, are in fact attributable to somatization rather than to depression per se. These findings have two important consequences. First, future research on depression should always control for the possible intervening effects of somatization. Second, the results of the present study show that the TRYCAT pathway could be a novel target in somatization. Attenuating the effects of IDO activation and in particular enhancing KAT or kynurenic acid levels could prove to be beneficial.

Conflicts of interest

The authors do not report any conflict of interest.

REFERENCES

- 1 Beck AT, Ward CH, Medelson M, Mock F, Erbaugh F (1961) An inventory for measuring depression. *Arch Gen Psychiatry* **4**: 561–571.
- 2 Bonaccorso S, Marino V, Puzella A, Pasquini M, Biondi M, Artini M, Almerighi C, Verkerk R, Meltzer H, Maes M (2002) Increased depressive ratings in patients with hepatitis C receiving interferon-alpha-based immunotherapy are related to interferon-alpha-induced changes in the serotonergic system. *J Clin Psychopharmacol* **22**(1): 86–90.
- 3 Booij L, Swenne CA, Brosschot JF, Haffmans PM, Thayer JF, Van der Does AJ (2006) Tryptophan depletion affects heart rate variability and impulsivity in remitted depressed patients with a history of suicidal ideation. *Biol Psychiatry* **60**(5): 507–514.
- 4 Brandacher G, Winkler C, Schroecksnadel K, Margreiter R, Fuchs D (2006) Antitumoral activity of interferon-gamma involved in impaired immune function in cancer patients. *Curr Drug Metab* **7**(6): 599–612.
- 5 Brkic S, Tomic S, Maric D, Novakov Mikic A, Turkulov V (2010). Lipid peroxidation is elevated in female patients with chronic fatigue syndrome. *Med Sci Monit*. **16**(12): CR628-32.
- 6 Brown FW, Golding JM, Smith GR Jr (1990) Psychiatric comorbidity in primary care somatization disorder. *Psychosom Med* **52**(4): 445–451.
- 7 Burdette SD, Parilo, MA, Kaplan LJ, Bailey H (2010) Systemic Inflammatory Response Syndrome. <http://emedicine.medscape.com/article/168943-overview>. Medscape, eMedicine.
- 8 Cairns BE, Svensson P, Wang K, Hupfeld S, Graven-Nielsen T, Sessle BJ, Berde CB, Arendt-Nielsen L (2003) Activation of peripheral NMDA receptors contributes to human pain and rat afferent discharges evoked by injection of glutamate into the masseter muscle. *J Neurophysiol* **90**(4): 2098–2105.
- 9 Cowen PJ, Parry-Billings M, Newsholme EA (1989) Decreased plasma tryptophan levels in major depression. *J Affect Disord* **16**(1): 27–31.
- 10 Curzon G (1979) Study of disturbed tryptophan metabolism in depressive illness. *Ann Biol Clin (Paris)* **37**(1): 27–33.
- 11 Curzon G, Sarna GS (1984) Tryptophan transport to the brain: newer findings and older ones reconsidered. In: Schlossberger HG, Kochen W, Linzen B, Steinhart H, editors. *Progress in Tryptophan and Serotonin Research*. Berlin: Walter De Gruyter, pp 145–157.

- 12 Davies SJ, Hood SD, Argyropoulos SV, Morris K, Bell C, Witchel HJ, Jackson PR, Nutt DJ, Potokar JP (2006) Depleting serotonin enhances both cardiovascular and psychological stress reactivity in recovered patients with anxiety disorders. *J Clin Psychopharmacol* **26**(4): 414–418.
- 13 DeMyer MK, Shea PA, Hendrie HC, Yoshimura NN (1981) Plasma tryptophan and five other amino acids in depressed and normal subjects. *Arch Gen Psychiatry* **38**(6): 642–646.
- 14 Derogatis LR (1994) Symptom Checklist-90-R. Administration, Scoring, and Procedures Manual, ed 3. Minneapolis: National Computer Systems.
- 15 Drummond PD (2006) Tryptophan depletion increases nausea, headache and photophobia in migraine sufferers. *Cephalalgia* **26**(10): 1225–1233.
- 16 Escobar Ji, Burnam MA, Karno M, Forsythe A, Golding JM (1987) Somatization in the community. *Arch Gen Psychiatry* **44**(8): 713–718.
- 17 Escobar Ji, Rubio-Stipec M, Canino G, Karno M (1989) Somatic symptoms index (SSI): A new and abridged somatization construct – Prevalence and epidemiological correlates in two large community samples. *J Nerv Ment Dis* **177**: 140–146.
- 18 Fernstrom JD (1984) Tryptophan availability and serotonin synthesis in rat brain: effects of experimental diabetes. In: Schlossberger HG, Kochen W, Linzen B, Steinhart H, editors. *Progress in Tryptophan and Serotonin Research*. Berlin: Walter de Gruyter, pp 161–172.
- 19 Gabbay V, Klein RG, Katz Y, Mendoza S, Guttman LE, Alonso CM, Babb JS, Hirsch GS, Liebes L (2010) The possible role of the kynurenine pathway in adolescent depression with melancholic features. *J Child Psychol Psychiatry* Apr 12.
- 20 Harvey JA, Schlosberg AJ, Yungler LM (1975) Behavioral correlates of serotonin depletion. *Fed Proc* **34**(9): 1796–1801.
- 21 Hervé C, Beyne P, Jamaalt H, Delacoux E (1996) Determination of tryptophan and its kynurenine pathway metabolites in human serum by high-performance liquid chromatography with simultaneous ultraviolet and fluorimetric detection. *J Chromatogr B Biomed Appl* **675**(1): 157–161.
- 22 Hiller W, Zaudig M, Mombour W (1997) *International Diagnostic Check Lists for ICD-10 and DSM-IV*. Bern: Huber.
- 23 Huensberg M, Winer J, Round R, Gompels M, Shahmanesh M (1998) Elevated serum kynurenine level is associated with peripheral neuropathy in HIV. *Neuroscience of HIV Infection. J Neurovirol* **4**: 353.
- 24 Joseph MS, Brewerton TD, Reus VI, Stebbins GT (1984) Plasma L-tryptophan/neutral amino acid ratio and dexamethasone suppression in depression. *Psychiatry Res* **11**(3): 185–192.
- 25 Kanai M, Funakoshi H, Takahashi H, Hayakawa T, Mizuno S, Matsumoto K, Nakamura T (2009) Tryptophan 2,3-dioxygenase is a key modulator of physiological neurogenesis and anxiety-related behavior in mice. *Mol Brain* **2**(1): 8.
- 26 Kaszaki J, Palásthy Z, Erczes D, Rácz A, Torday C, Varga G, Vécsei L, Boros M (2008) Kynurenic acid inhibits intestinal hypermotility and xanthine oxidase activity during experimental colon obstruction in dogs. *Neurogastroenterol Motil* **20**(1): 53–62.
- 27 Kékesi G, Horváth G (2002) Analgesic effect of kynurenic acid. *Ideggyogy Sz* **55**(9–10): 313–322.
- 28 Kilkens TO, Honig A, van Nieuwenhoven MA, Riedel WJ, Brummer RJ (2004) Acute tryptophan depletion affects brain-gut responses in irritable bowel syndrome patients and controls. *Gut* **53**(12): 1794–1800.
- 29 Kroenke K, Spitzer RL, deGruy FV 3rd, Hahn SR, Linzer M, Williams JB, Brody D, Davies M (1997) Multisomatoform disorder. An alternative to undifferentiated somatoform disorder for the somatizing patient in primary care. *Arch Gen Psychiatry* **54**(4): 352–358.
- 30 Ladwig KH, Marten-Mittag B, Erazo N, Gündel H (2001) Identifying somatization disorder in a population-based health examination survey: psychosocial burden and gender differences. *Psychosomatics* **42**(6): 511–518.
- 31 Lehmann J (1972) Mental and neuromuscular symptoms in tryptophan deficiency: a review with special reference to mental symptoms in a selected material of carcinoid patients. *Acta Psychiatr Scand (Suppl)* 237.
- 32 Li J, McRoberts JA, Ennes HS, Trevisani M, Nicoletti P, Mittal Y, Mayer EA (2006) Experimental colitis modulates the functional properties of NMDA receptors in dorsal root ganglia neurons. *Am J Physiol Gastrointest Liver Physiol* **291**(2): G219–228.
- 33 Maes M (1995) Evidence for an immune response in major depression: a review and hypothesis. *Prog Neuropsychopharmacol Biol Psychiatr* **19**(1): 11–38.
- 34 Maes M (2010) Depression is an inflammatory disease, but cell-mediated immune activation is the key component of depression. *Prog Neuropsychopharmacol Biol Psychiatry* [Epub ahead of print] PubMed PMID: 20599581.
- 35 Maes M, Meltzer HY (1995) The Serotonin Hypothesis of major depression. Selected Chapters on Mood Disorders. In: F.Bloom, D.Kupfer, editors. *Psychopharmacology, The Fourth Generation of Progress*. New York: Raven Press, pp 933–944.
- 36 Maes M, Twisk FN (2010). Chronic fatigue syndrome: Harvey and Wessely's (bio)psychosocial model versus a bio(psychosocial) model based on inflammatory and oxidative and nitrosative stress pathways. *BMC Med.* **8**: 35. Review.
- 37 Maes MH, De Ruyter M, Suy E (1987a) Prediction of subtype and severity of depression by means of dexamethasone suppression test, L-tryptophan: competing amino acid ratio, and MHPG flow. *Biol Psychiatry* **22**(2): 177–188.
- 38 Maes M, De Ruyter M, Suy E (1987b) The renal excretion of xanthurenic acid following L-tryptophan loading in depressed patients. *Human Psychopharmacol Clin Exp* **2**(4): 231–235.
- 39 Maes M, Jacobs MP, Suy E, Vandewoude M, Minner B, Raus J (1990a) Effects of dexamethasone on the availability of L-tryptophan and on the insulin and FFA concentrations in unipolar depressed patients. *Biol Psychiatry* **27**(8): 854–862.
- 40 Maes M, Maes L, Suy E (1990b) Symptom profiles of biological markers in depression: a multivariate study. *Psychoneuroendocrinology* **15**(1): 29–37.
- 41 Maes M, Schotte C, Scharpé S, Martin M, Blockx P (1990c) The effects of glucocorticoids on the availability of L-tryptophan and tyrosine in the plasma of depressed patients. *J Affect Disord* **18**(2): 121–127.
- 42 Maes M, Minner B, Suy E (1991a). The relationships between the availability of L-tryptophan to the brain, the spontaneous HPA-axis activity, and the HPA-axis responses to dexamethasone in depressed patients. *J AminoAcids* **1**(1): 57–65
- 43 Maes M, Vandewoude M, Scharpé S, De Clercq L, Stevens W, Lepoutre L, Schotte C (1991b) Anthropometric and biochemical assessment of the nutritional state in depression: evidence for lower visceral protein plasma levels in depression. *J Affect Disord* **23**(1): 25–33.
- 44 Maes M, Meltzer HY, Scharpé S, Bosmans E, Suy E, De Meester I, Calabrese J, Cosyns P (1993) Relationships between lower plasma L-tryptophan levels and immune-inflammatory variables in depression. *Psychiatry Res* **49**(2): 151–165.
- 45 Maes M, Scharpé S, Meltzer HY, Okayli G, Bosmans E, D'Hondt P, Vanden Bossche BV, Cosyns P (1994) Increased neopterin and interferon-gamma secretion and lower availability of L-tryptophan in major depression: further evidence for an immune response. *Psychiatry Res* **54**(2): 143–160.
- 46 Maes M, Wauters A, Verkerk R, Demedts P, Neels H, Van Gastel A, Cosyns P, Scharpé S, Desnyder R (1996) Lower serum L-tryptophan availability in depression as a marker of a more generalized disorder in protein metabolism. *Neuropsychopharmacology* **15**(3): 243–251.
- 47 Maes M, Verkerk R, Vandoolaeghe E, Van Hunsel F, Neels H, Wauters A, Demedts P, Scharpé S (1997) Serotonin-immune interactions in major depression: lower serum tryptophan as a marker of an immune-inflammatory response. *Eur Arch Psychiatry Clin Neurosci* **247**(3): 154–161.
- 48 Maes M, Bonaccorso S, Marino V, Puzella A, Pasquini M, Biondi M, Artini M, Almerighi C, Meltzer H (2001a) Treatment with interferon-alpha (IFN alpha) of hepatitis C patients induces lower serum dipeptidyl peptidase IV activity, which is related to IFN alpha-induced depressive and anxiety symptoms and immune activation. *Mol Psychiatry* **6**(4): 475–480.
- 49 Maes M, Ombelet W, Verkerk R, Bosmans E, Scharpé S (2001b) Effects of pregnancy and delivery on the availability of plasma tryptophan to the brain: relationships to delivery-induced immune activation and early post-partum anxiety and depression. *Psychol Med* **31**(5): 847–858.

- 50 Maes M, Verkerk R, Bonaccorso S, Ombet W, Bosmans E, Scharpé S (2002) Depressive and anxiety symptoms in the early puerperium are related to increased degradation of tryptophan into kynurenine, a phenomenon which is related to immune activation. *Life Sci* **71**(16): 1837–1848
- 51 Maes M, Mihaylova I, Ruyter MD, Kubera M, Bosmans E (2007) The immune effects of TRYCATs (tryptophan catabolites along the IDO pathway): relevance for depression – and other conditions characterized by tryptophan depletion induced by inflammation. *Neuro Endocrinol Lett* **28**(6): 826–831.
- 52 Maes M, Leonard BE, Myint AM, Kubera M, Verkerk R (2010) The new 5-HT hypothesis of depression: cell-mediated immune activation induces indoleamine 2,3-dioxygenase, which leads to lower plasma tryptophan and synthesis of detrimental tryptophan catabolites (TRYCATs), both of which contribute to the onset of depression. *Prog Neuropsychopharmacol Biol Psychiatry* accepted.
- 53 Maes M, Kubera M, Uytterhoeven M, Vrydags N, Bosmans E (2011). Increased plasma peroxides as a marker of oxidative stress in myalgic encephalomyelitis/chronic fatigue syndrome (ME/CF). *Med Sci Monit*. **17**(4): SC11-5.
- 54 McRoberts JA, Coutinho SV, Marvizón JC, Grady EF, Tognetto M, Sengupta JN, Ennes HS, Chaban VV, Amadesi S, Creminon C, Lanthorn T, Geppetti P, Bunnett NW, Mayer EA (2001) Role of peripheral N-methyl-D-aspartate (NMDA) receptors in visceral nociception in rats. *Gastroenterology* **120**(7): 1737–1748.
- 55 Mees L, Tuboly G, Nagy E, Benedek G, Horvath G (2009) The peripheral antinociceptive effects of endomorphin-1 and kynurenic acid in the rat inflamed joint model. *Anesth Analg* **109**(4): 1297–1304.
- 56 Moir AT, Eccleston D (1968) The effects of precursor loading in the cerebral metabolism of 5-hydroxyindoles. *J Neurochem* **15**(10): 1093–1108.
- 57 Møller SE (1985) Tryptophan to competing amino acids ratio in depressive disorder: relation to efficacy of antidepressive treatments. *Acta Psychiatr Scand Suppl* **325**: 3–31.
- 58 Morgan CJ, Badawy AA (1989) Effects of a suppression test dose of dexamethasone on tryptophan metabolism and disposition in the rat. *Biol Psychiatry* **25**(3): 359–362.
- 59 Moroni F, Russi P, Gallo-Mezo MA, Moneti G, Pellicciari R (1991) Modulation of quinolinic and kynurenic acid content in the rat brain: effects of endotoxins and nicotinylalanine. *J Neurochem* **57**: 1630–1635.
- 60 Myint AM, Kim YK, Verkerk R, Scharpé S, Steinbusch H, Leonard B (2007) Kynurenine pathway in major depression: evidence of impaired neuroprotection. *J Affect Disord* **98**(1–2): 143–151.
- 61 Németh H, Toldi J, Vécsei L (2005) Role of kynurenines in the central and peripheral nervous systems. *Curr Neurovasc Res* **2**(3): 249–260.
- 62 Oxenkrug GF (2007) Genetic and hormonal regulation of tryptophan kynurenine metabolism: implications for vascular cognitive impairment, major depressive disorder, and aging. *Ann N Y Acad Sci* **1122**: 35–49.
- 63 Raison CL, Dantzer R, Kelley KW, Lawson MA, Woolwine BJ, Vogt G, Spivey JR, Saito K, Miller AH (2010) CSF concentrations of brain tryptophan and kynurenines during immune stimulation with IFN- α : relationship to CNS immune responses and depression. *Mol Psychiatry* **15**(4): 393–403.
- 64 Richell RA, Deakin JF, Anderson IM (2005) Effect of acute tryptophan depletion on the response to controllable and uncontrollable noise stress. *Biol Psychiatry* **57**(3): 295–300.
- 65 Rief W, Fichter M (1992) The Symptom Check List SCL-90-R and its ability to discriminate between dysthymia, anxiety disorders, and anorexia nervosa. *Psychopathology* **25**(3): 128–138.
- 66 Rief W, Hiller W, Heuser J (1997) SOMS – Das Screening für somatoforme Störungen. Manual zum Fragebogen. Bern: Huber.
- 67 Rief W, Hessel A, Braehler E (2001a) Somatization symptoms and hypochondriacal features in the general population. *Psychosom Med* **63**(4): 595–602.
- 68 Rief W, Pilger F, Ihle D, Bosmans E, Egyed B, Maes M (2001b) Immunological differences between patients with major depression and somatization syndrome. *Psychiatry Res* **105**(3): 165–174.
- 69 Rief W, Pilger F, Ihle D, Verkerk R, Scharpe S, Maes M (2004) Psychobiological aspects of somatoform disorders: contributions of monoaminergic transmitter systems. *Neuropsychobiology* **49**(1): 24–29.
- 70 Rief W, Hennings A, Riemer S, Euteneuer F (2010) Psychobiological differences between depression and somatization. *J Psychosom Res* **68**(5): 495–502.
- 71 Simon GE, VonKorff M (1991) Somatization and psychiatric disorder in the NIMH Epidemiologic Catchment Area study. *Am J Psychiatry* **148**(11): 1494–1500.
- 72 Song C, Dinan T, Leonard BE (1994) Changes in immunoglobulin, complement and acute phase protein levels in the depressed patients and normal controls. *J Affect Disord* **30**(4): 283–288.
- 73 Song C, Lin A, Bonaccorso S, Heide C, Verkerk R, Kenis G, Bosmans E, Scharpe S, Whelan A, Cosyns P, de Jongh R, Maes M (1998) The inflammatory response system and the availability of plasma tryptophan in patients with primary sleep disorders and major depression. *J Affect Disord* **49**(3): 211–219.
- 74 Stone TW, Darlington LG (2002) Endogenous kynurenines as targets for drug discovery and development. *Nat Rev Drug Discov* **1**(8): 609–620.
- 75 Struzik L, Duffin J, Vermani M, Hegadoren K, Katzman MA (2002) Effects of tryptophan depletion on central and peripheral chemoreflexes in man. *Respir Physiol Neurobiol* **133**(3): 183–195.
- 76 Swardfager W, Herrmann N, Dowlati Y, Oh PI, Kiss A, Walker SE, Lanctôt KL (2009) Indoleamine 2,3-dioxygenase activation and depressive symptoms in patients with coronary artery disease. *Psychoneuroendocrinology* **34**(10): 1560–1566.
- 77 Swartz M, Landerman R, George L, Blazer D, Escobar J (1990) Somatization disorder. In: Robins LN, Regier D, editors. *Psychiatric Disorders in America*. New York, Free Press.
- 78 Takikawa O, Yoshida R, Yasui H, Hayaishi O (1984) The relationship between plasma kynurenine and indoleamine 2,3-dioxygenase activity in the extrahepatic tissues. In: Schlossberger HG, Kochen W, Linzen B, Steinhart H, editors. *Progress in Tryptophan and Serotonin Research*. Berlin-New York: Walter de Gruyter and Co., pp 517–520.
- 79 Tanke MA, Alserda E, Doornbos B, van der Most PJ, Goeman K, Postema F, Korf J (2008) Low tryptophan diet increases stress-sensitivity, but does not affect habituation in rats. *Neurochem Int* **52**(1–2): 272–281.
- 80 Terre L, Poston WS, Foreyt J, St Jeor ST (2003) Do somatic complaints predict subsequent symptoms of depression? *Psychother Psychosom* **72**(5): 261–267.
- 81 van Veen JF, van Vliet IM, de Rijk RH, van Pelt J, Mertens B, Fekkes D, Zitman FG (2009) Tryptophan depletion affects the autonomic stress response in generalized social anxiety disorder. *Psychoneuroendocrinology* **34**(10): 1590–1594.
- 82 Wichers MC, Koek GH, Robaey G, Praamstra AJ, Maes M (2005a) Early increase in vegetative symptoms predicts IFN- α -induced cognitive-depressive changes. *Psychol Med* **35**(3): 433–441.
- 83 Wichers MC, Koek GH, Robaey G, Verkerk R, Scharpé S, Maes M (2005b) IDO and interferon- α -induced depressive symptoms: a shift in hypothesis from tryptophan depletion to neurotoxicity. *Mol Psychiatry* **10**(6): 538–544.
- 84 Wittchen HU, Schramm E, Zaudig M, Unland H (1997) SKID: Stukturiertes klinisches Interview für DSM-IV, Achse I, deutsche Version. Göttingen: Hogrefe.
- 85 Wu LJ, Zhuo M (2009) Targeting the NMDA receptor subunit NR2B for the treatment of neuropathic pain. *Neurotherapeutics* **6**(4): 693–702.
- 86 Yang X, Yang HB, Xie QJ, Liu XH, Hu XD (2009) Peripheral inflammation increased the synaptic expression of NMDA receptors in spinal dorsal horn. *Pain* **144**(1–2): 162–169.
- 87 Young SN (1981) Mechanism of decline in rat brain 5-hydroxytryptamine after induction of liver tryptophan pyrrolase by hydrocortisone: roles of tryptophan catabolism and kynurenine synthesis. *Br J Pharmacol* **74**(3): 695–700.
- 88 Yuwiler A, Oldendorf WH, Geller E, Braun L (1977) Effect of albumin binding and amino acid competition on tryptophan uptake into brain. *J Neurochem* **28**(5): 1015–1023.
- 89 Zepf FD, Holtmann M, Stadler C, Wöckel L, Poustka F (2009) Reduced serotonergic functioning changes heart rate in ADHD. *J Neural Transm* **116**(1): 105–108.